

北太平洋における東方伝播する十年規模貯熱量シグナルの起源 Origin of Decadal-scale Eastward-propagating Signals of Oceanic Heat Content in the North Pacific Ocean

田口 文明^{1*}, ニクラス シュナイダー²
TAGUCHI, Bunmei^{1*}, Niklas Schneider²

¹ 海洋研究開発機構 地球シミュレータセンター, ² ハワイ大学 国際太平洋研究センター

¹Japan Agency for Marine-Earth Science and Technology, ²International Pacific Research Center, University of Hawaii at Manoa

The mid-latitude North Pacific Ocean supports both eastward and westward moving subsurface signals. Typically studies of sea surface height anomalies report westward propagation and interpret these as the first baroclinic mode Rossby waves (RWs), while investigations of upper ocean heat content (OHC) find eastward propagation and cite mean flow advection as an underlying cause. Yet, OHC in general affects the velocity field via density change and thus also evolves as RWs. We investigate this dichotomy using a 150-year CGCM integration. Simulated OHC signals are distinguished in terms of two processes that can support eastward propagation: higher baroclinic RW modes and density-compensated temperature and salinity anomalies, a.k.a. spiciness. Our analysis suggests a unique role of the Kuroshio/Oyashio Extension (KOE) region as an origin of the spiciness and higher mode RW signals as follows. First, wind-forced, westward-propagating first baroclinic RW causes circulation anomalies in the KOE region, accompanying the meridional shift of the subarctic front. Anomalous advection of mean temperature and salinity gradients then generates spiciness anomalies, which are advected eastward by mean currents, manifested as eastward co-propagation of OHC and sea surface salinity anomalies. While being advected, the surface temperature anomaly associated with the spiciness signal is damped by air-sea heat exchange and thus generates density anomalies, which further propagate eastward as higher mode RWs. The result suggests that the commonly used indices of OHC of the upper ocean result from a mixture of different dynamics that are transformed one another in the KOE region.

キーワード: 海洋表層貯熱量, 十年規模変動, ロスビー波, 密度補償シグナル, 大気海洋相互作用, 亜寒帯前線

Keywords: Oceanic Heat Content, Decadal Variability, Rossby Waves, Spiciness, Air-sea Interaction, Subarctic Frontal Zone